

Intrazelluläre Borrelien – Stress (bzw. stealth-) Varianten Stay of Borrelia stress – variants in the cells of the host organism

Intrazelluläre Lebensformen können nicht mit Antibiotika erreicht werden, wenn sie sich monatelang nicht vermehren.

Intracellular life forms cannot be reached with antibiotics when they do not reproduce for months.

Valesova, M., et al., (1989) Detection of Borrelia in the synovial tissue from a patient with Lyme borreliosis by electron microscopy. J Rheumatol, 16(11), 1502-5.

Stanek, G., et al.(1990) Isolation of Borrelia burgdorferi from the myocardium of a patient with longstanding cardiomyopathy. N Engl J Med, 322(4), 249-52

Ma, Y., Sturrock A, Weis J (1991) **Intracellular** localization of Borrelia burgdorferi within human endothelial cells. Infect Immun, 59(2), 671-8

Georgilis K, Peacocke M, Klempner MS.(1992) Fibroblasts protect the Lyme disease spirochete, Borrelia burgdorferi, from ceftriaxone in vitro. J Infect Dis.166(2), 440- 4.

Klempner MS, Noring R, Rogers RA.(1993) Invasion of human skin fibroblasts by the Lyme disease spirochete, Borrelia burgdorferi. J Infect Dis. 167(5), 1074- 81.

Haupt, T., et al., (1993) Persistence of Borrelia burgdorferi in ligamentous tissue from a patient with chronic Lyme borreliosis. Arthritis Rheum, 36(11), 1621-6.

Hulinska D, Basta J, Murgia R, Cinco M (1995) **Intracellular** morphological events observed by electron microscopy on neutrophil phagocytosis of Borrelia garinii Journal of Spirochetal and Tick-Borne Diseases. 2(4), 82-86.

Girschick HJ, Huppertz HI, Russmann H, Krenn V, Karch H. (1996) **Intracellular** persistence of Borrelia burgdorferi in human synovial cells. Rheumatol Int 16(3), 125–132

Aberer, E. et al. (1996) Heterogeneity of Borrelia burgdorferi in the skin. Am J Dermatopathol, 18(6), 571-9

Brouqui P, Badiaga S, and Raoult D (1996). ["Eucaryotic cells protect Borrelia burgdorferi from the action of penicillin and ceftriaxone but not from the action of doxycycline and erythromycin"](#) (PDF). Antimicrob Agents Chemother 40 (6), 1552–1554.

Nanagara, R., P.H. Duray, and H.R. Schumacher, Jr. (1996) Ultrastructural demonstration of spirochetal antigens in synovial fluid and synovial membrane in chronic Lyme disease: possible factors contributing to persistence of organisms. Hum Pathol, 27(10), 1025-34.

Dorward, D.W., E.R. Fischer, D.M. Brooks, (1997) Invasion and cytopathic killing of human lymphocytes by spirochetes causing Lyme disease. Clin Infect Dis, 25 Suppl 1, 2-8

Priem, S., et al.(1998) Detection of Borrelia burgdorferi by polymerase chain reaction in synovial membrane, but not in synovial fluid from patients with persisting Lyme arthritis after antibiotic therapy. Ann Rheum Dis, 57(2), 118-21

de Koning, J., et al. (1998) Demonstration of spirochetes in cardiac biopsies of patients with Lyme disease. J Infect Dis, 160(1), 150-3

Chary-Valckenaere I, Jaulhac B, Champigneulle J, et al. (1998) Ultrastructural demonstration of intracellular localization of *Borrelia burgdorferi* in Lyme arthritis. *Br J Rheumatol* 37, 468-470.

Kroun M (2003) TNF and interruption of apoptosis (programmed cell death) seem to play an important role in persistence of most (all?) chronic INTRACELLULAR INFECTIONS.

<http://lymerick.net/NFkB.htm>

Duray, P.H., et al. (2005) Invasion of human tissue **ex vivo** by *Borrelia burgdorferi*. *J Infect Dis*, 191(10), 1747-54.

Livengood, J.A. and R.D. Gilmore, Jr., (2006) Invasion of human neuronal and glial cells by an infectious strain of *Borrelia burgdorferi*. *Microbes Infect*, 8(14-15), 2832-40

Margulis L, Maniotis A, MacAllister J et al. (2009) Position paper. Spirochete round bodies Syphilis, Lyme disease & AIDS: Resurgence of „the great imitator“?. *Symbiosis* 47, 51-58

<http://www1.biogema.de/WEK/312-Margulis-final.pdf>

Müller, KE (2009) Erkrankung der elastischen und kollagenen Fasern von Haut, Sehnen und Bändern bei Lyme-Borreliose. *Umwelt Medizin Gesellschaft* 22, 112-118.

<http://www.schattenblick.de/infopool/medizin/fachmed/mz1um194.html>

Chmielewski T, Tylewska-Wierzhanowska S (2011) Inhibition of fibroblast apoptosis by *Borrelia afzelii*, *Coxiella burnetii* and *Bartonella henselae*. *Poll Microbiol* 60(3). 269-72

[Asner SA](#), [Morré SA](#), [Bochud PY](#), [Greub G](#). (2014) **Host factors and genetic susceptibility for infections due to intracellular bacteria and fastidious organisms.** *Clin Microbiol Infect*. doi:

10.1111/1469-0691.12806. [Epub ahead of print] <http://www.ncbi.nlm.nih.gov/pubmed/25366416>

„In conclusion, this review supports the paramount role of genetic factors in clinical presentations and severity of infections caused by intracellular fastidious bacteria. Genetic predisposition should be further explored with more performing techniques such as exome sequencing ».

- ➔ **Bakterien Stressvarianten** <http://www.erlebnishaft.de/stressvar1.pdf>
- ➔ **Borrelien – Populationsdynamik, Stressvarianten** <http://www.erlebnishaft.de/stressvar2.pdf>

- ➔ **Genetische Faktoren** http://www.xerlebnishaft.de/genetische_faktoren.pdf
http://www.erlebnishaft.de/TLR2_1_3_7_13.pdf

[Bernt - Dieter Huismans](#), Start 2013. Letzte Revision Februar 2017 www.Huismans.click



Back to top: http://www.xerlebnishaft.de/borr_intrazellulaer.pdf